

KUNS-1509
 HE(TH) 98/7
 UT-813
 hep-th/9805119

On the Generalized Gluing and Resmoothing Theorem

Tsuguhiko ASAKAWA,^{*)} Taichiro KUGO^{**)} and Tomohiko TAKAHASHI^{†,***)}

Department of Physics, Kyoto University, Kyoto 606-01

[†]*Department of Physics, University of Tokyo, Tokyo 113*

Abstract

The generalized gluing and resmoothing theorem originally proved by LeClair, Peskin and Preitschopf, gives a powerful formula for the fused vertex obtained by contracting any two vertices in string field theories. Although the theorem is naturally expected to hold for the vertices at any loop level, the original proof was restricted to the vertices at tree level. Here we present a simplified proof for the tree level theorem and then prove explicitly the extended version at one-loop level. We also find that a non-trivial sign factor, which depends on the string states to be contracted, appears in the theorem. This sign factor turns out to be essential for reproducing correctly the conformal field theory correlation function on the torus.

^{*)} E-mail: asakawa@gauge.scphys.kyoto-u.ac.jp

^{**)} E-mail: kugo@gauge.scphys.kyoto-u.ac.jp

^{***)} JSPS Research Fellow. E-mail: tomo@hep-th.phys.s.u-tokyo.ac.jp

§1. Introduction

The basic ingredients of string field theories (SFT) are the vertices. We need to combine (or fuse) the various vertices in many situations, e.g., in showing the gauge invariance of the SFT action and in computing string scattering amplitudes in perturbation theory. LeClair, Peskin and Preitschopf¹⁾ (LPP) have developed a powerful method for defining general multistring vertices by conformally mapping the unit disks of participating strings into a complex plane and by using the correlation functions of conformal field theory^{3), 4)} (CFT) in the plane. They then proved a theorem, which they called “Generalized Gluing and Resmoothing Theorem” (GGRT),²⁾ giving a general formula for the fused vertex obtained by a contraction of two vertices. They showed that the fused vertex is just equals their multistring vertex corresponding to the conformal mappings induced by gluing the two world sheets into one.

The point here is that the equality holds with *weight one* if the conformal anomaly (i.e., central charge c) is zero. Their proof is very thorough and even pedagogical. It is, however, a bit complicated and they did not take much care about the sign of the equality. The relevant sign factor is not a mere phase but an operator which changes sign depending on the string states to be contracted, and become very important, for example, for the cancellations of divergences between two graphs⁵⁾ and also in the proof of the gauge invariance of the SFT action.⁶⁾ In the context of this paper also, the sign factor in the theorem at tree level turns out to be essential for reproducing correctly the conformal field theory correlation function on the torus.

Here in this paper we first present a much simplified proof for the LPP GGRT, and determine the sign of the equality carefully. Our proof is inspired by the sewing method of two conformal field theories defined on two Riemann surfaces, which has been given, in particular, by Sonoda.⁷⁾ Actually, the GGRT by LPP is a SFT version of this general way of sewing two CFT’s. The gluing and the sewing are essentially the same and are just the insertion of the complete set of states. So, although the original GGRT by LPP is restricted to the vertices at tree level, it is naturally expected to hold at any loop level. Nevertheless, the SFT version is not so trivial. This is because the gluing in SFT must be performed by contracting two strings, one each from the two vertices. On the other hand the sewing of two CFT’s is performed by excising two holes freely, one each on the two Riemann surfaces. To do the same thing in the SFT case and to make contact with the definition of the vertices, one needs to map the string world sheets back and forth. These mappings give non-trivial conformal transformations on the operators, which must be traced neatly. We perform this procedure and prove explicitly an extended version of the GGRT at one loop level. It may

be interesting to note that, as a byproduct, the formula for the CFT correlation function on the torus⁸⁾

$$\langle \mathcal{O}_1 \mathcal{O}_2 \cdots \rangle_{\text{torus } \tau} = \text{Tr} [(-1)^{N_{\text{FP}}} q^{2L_0} \mathcal{O}_1 \mathcal{O}_2 \cdots] \quad (1.1)$$

is automatically *derived* by this procedure (up to an overall factor convention), where $q = e^{i\pi\tau}$ and N_{FP} is the ghost number operator. The factor $(-1)^{N_{\text{FP}}}$ in this expression comes from the operator sign factor in the theorem at tree level mentioned above.

This paper is organized as follows. First, in §2, we briefly review the the definition of the vertices and the GGRT given by LPP. In §3, after making some remarks on the ambiguity present when $c \neq 0$ in defining conformal transformation operators U_f corresponding to the mappings $f(z)$, we present two propositions to clarify when U_f leaves inert the $\text{SL}(2;\mathbb{C})$ bra and ket vacua, and then, gives a simplified proof for the GGRT of LPP. The extension of GGRT to one-loop level is given and proved in §4.

For simplicity of presentation, we assume henceforth that the strings are all bosonic open strings, so that the relevant conformal fields $\phi(z)$ are string coordinates $\partial X^\mu(z)$, and reparameterization ghost $c(z)$ and anti-ghost $b(z)$, possessing dimensions $d = 1, -1$ and 2 , respectively. Closed string can be treated similarly since it is more or less equivalent with a pair of open strings.

§2. GGRT at tree level

First of all, let us recall LPP's definition of the tree level vertex which refers to the conformal field theory in the complex plane (two dimensional manifold M which is topologically equal to S^2):^{1), 9), 10)}

$$\langle v(n, \cdots, 2, 1) | |A_1\rangle_1 |A_2\rangle_2 \cdots |A_n\rangle_n \equiv \left\langle h_1[\mathcal{O}_{A_1}] h_2[\mathcal{O}_{A_2}] \cdots h_n[\mathcal{O}_{A_n}] \right\rangle_M. \quad (2.1)$$

Here $\langle v(n, \cdots, 2, 1) |$ is the n -point LPP vertex, which is defined as a bra state in the product space $\otimes_{i=1}^n \mathcal{H}_i$ of n string Fock spaces $\mathcal{H}_i$, and each string state $|A_i\rangle_i \in \mathcal{H}_i$ are given in the form

$$|A\rangle_i = \mathcal{O}_A |0\rangle_i, \quad (2.2)$$

where $\mathcal{O}_A$ is an operator creating the state A of string i from the $\text{SL}(2;\mathbb{C})$ invariant vacuum $|0\rangle_i$ in $\mathcal{H}_i$; for instance, the tachyon state of momentum p is given by the vertex operator $\mathcal{O}(w) = c(w) \exp(ip \cdot X(w))$ at $w = 0$, and the ladder operators $\phi_n = \{\alpha_n, c_n, b_n\}$ are given by the contour integration $\oint (dw/2\pi i) w^{n+d-1} \phi(w)$ encircling the origin.

The meaning of the right-hand side of Eq. (2.1) is as follows: any vertex $\langle v(n, \cdots, 2, 1) |$ is defined by specifying how the participating strings i are glued each other. We can regard

each string world sheet from infinite past ($\tau_i = -\infty$) to the interaction time ($\tau_i = 0$) as a unit disk $|w_i| \leq 1$ with $w_i = \exp(\tau_i + i\sigma_i)$, and the world sheet formed by gluing those string sheets as a complex z -plane, which we call M ($\sim S^2$), for tree level vertex case. So this gluing can be simply specified by giving conformal mappings $h_i(w_i)$ of each string w_i plane into the complex z plane M , which is analytic and invertible inside the each unit circle $|w_i| = 1$. Generally, any conformal mapping $f: w \rightarrow z = f(w)$ of w -plane to z -plane also defines a mapping of operators $\mathcal{O}$ in the w -plane to operators $f[\mathcal{O}]$ in the z -plane:

$$f[\mathcal{O}] \equiv U_f \mathcal{O} U_f^{-1}. \quad (2.3)$$

If the operator $\mathcal{O}$ is a primary conformal field $\phi(w)$ of dimension d_ϕ , this mapping is defined to be

$$f[\phi(w)] = \left(\frac{df(w)}{dw} \right)^{d_\phi} \phi(f(w)). \quad (2.4)$$

The operator representation U_f in Eq. (2.3) of the conformal mapping f is uniquely determined by this transformation law (2.4) of the primary fields *up to a multiplicative constant*. Since the Fourier components of the energy momentum tensor $T(z)$, $L_n \equiv \oint (dz/2\pi i) z^{n+1} T(z)$, generate infinitesimal conformal transformations, the operator U_f for the finite transformation f can be given in the form

$$U_f = \exp\left(\sum_n v_{-n} L_n\right) \quad (2.5)$$

with certain parameters v_n . ($v(z) \equiv \sum_n v_{-n} z^{n+1}$ and $f(z)$ are related by $f(z) = e^{v(z)\partial_z} z$.) We should keep in mind, however, that this parameterization (2.5) for U_f is not unique and that the very definition of U_f by Eq. (2.3) has an ambiguity of multiplicative constant. We shall come back to this problem later in the next section.

Now the meaning of the right-hand side of Eq. (2.1) will be clear: it gives a correlation function of the mapped operators $h_i[\mathcal{O}_{A_i}]$ of the conformal field theory in the z plane. Note the crucial conceptual difference between both sides of the defining equation (2.1) of LPP vertex; the left-hand side is an inner product in the product space $\otimes_{i=1}^n \mathcal{H}_i$ of n string Fock spaces $\mathcal{H}_i$, while the right-hand side is a correlation function of a *single string* conformal field theory in z plane.

We need a little more preparation to state the GGRT. Let us introduce bra and ket reflectors $\langle R(1,2)|$ and $|R(1,2)\rangle$ which convert ket string states $|A\rangle$ to bra states $\langle A|$, and vice versa:

$$\langle R(1,2)| |A\rangle_2 = {}_1\langle A|, \quad {}_2\langle A| |R(2,1)\rangle = |A\rangle_1. \quad (2.6)$$

The reflectors $\langle R(1,2)|$ and $|R(1,2)\rangle$ are just the metric g_{IJ} and g^{IJ} , respectively, if we use notation $|A\rangle \equiv A^I$ and $\langle A| \equiv A_I$. So they can be defined by giving an inner product in the

string Fock space $\mathcal{H}$. A natural inner product³⁾ is defined by using the inversion $I(z) = -1/z$ as follows:

$$\begin{aligned}\langle A|B\rangle &= \langle R(1,2)| |A\rangle_2 |B\rangle_1 \\ &= \langle I[\mathcal{O}_A] \mathcal{O}_B \rangle .\end{aligned}\tag{2.7}$$

It is easy to find an explicit oscillator expression for the reflectors, as can be found, e.g., in Refs. 2) and 11). We here need not that explicit expression but the following formal one. Let $\{|\alpha\rangle\}$ be a complete set of the ket string states and $\{|\tilde{\alpha}\rangle\}$ be its orthonormal dual under this inner product; i.e., $\langle \tilde{\beta}|\alpha\rangle = \langle I[\mathcal{O}_{\tilde{\beta}}] \mathcal{O}_{\alpha} \rangle = \delta_{\beta}^{\alpha}$. Then we have a completeness relation:

$$\sum_{\alpha} |\alpha\rangle \langle \tilde{\alpha}| = \sum_{\alpha} \mathcal{O}_{\alpha} |0\rangle \langle 0| I[\mathcal{O}_{\tilde{\alpha}}] = \mathbf{1} \quad \text{in } \mathcal{H},\tag{2.8}$$

where $\mathcal{O}_{\alpha}$ and $\mathcal{O}_{\tilde{\alpha}}$ are operators creating the states $|\alpha\rangle$ and $|\tilde{\alpha}\rangle$, respectively. It is now clear that the reflectors have the following formal expressions:

$$\begin{aligned}\langle R(1,2)| &= \sum_{\alpha} {}_1\langle \alpha| {}_2\langle \tilde{\alpha}| = \sum_{\alpha} {}_1\langle 0| I[\mathcal{O}_{\alpha}] {}_2\langle 0| I[\mathcal{O}_{\tilde{\alpha}}], \\ |R(1,2)\rangle &= \sum_{\alpha} |\alpha\rangle_1 |\tilde{\alpha}\rangle_2 = \sum_{\alpha} \mathcal{O}_{\alpha} |0\rangle_1 \mathcal{O}_{\tilde{\alpha}} |0\rangle_2 .\end{aligned}\tag{2.9}$$

Now we define the contraction (or fusion) of two vertices appearing in the GGRT. Let $\langle v(C, \{A_i\})|$ be an $(n+1)$ -point LPP vertex for the strings A_i ($i = 1, 2, \dots, n$) and C defined by conformal mappings h_{A_i} and h_C :

$$\langle v(C, \{A_i\})| \prod_{i=1}^n |A_i\rangle_{A_i} |C\rangle_C \equiv \left\langle \prod_{i=1}^n h_{A_i}[\mathcal{O}_{A_i}] h_C[\mathcal{O}_C] \right\rangle_M .\tag{2.10}$$

And let $\langle v(D, \{B_j\})|$ be another $(m+1)$ -point LPP vertex defined similarly:

$$\langle v(D, \{B_j\})| \prod_{j=1}^m |B_j\rangle_{B_j} |D\rangle_D \equiv \left\langle \prod_{j=1}^m h_{B_j}[\mathcal{O}_{B_j}] h_D[\mathcal{O}_D] \right\rangle_N .\tag{2.11}$$

Note that we have called the z -planes for the two cases M and N , for distinction, although they are both topologically $\sim S^2$. Then we can define a fused vertex $\langle v(\{B_j\}, \{A_i\})|_{\text{fused}}$ of these two vertices by gluing the strings C and D in each by the help of the ket reflector $|R(C, D)\rangle$:

$$\langle v(\{B_j\}, \{A_i\})|_{\text{fused}} \equiv \langle v(D, \{B_j\})| \langle v(C, \{A_i\})| |R(C, D)\rangle .\tag{2.12}$$

Intuitively, the fusion gives a new Riemann surface which is formed by cutting out the images of the unit disks of strings C and D in M and N , respectively, and then gluing smoothly the rest pieces of M and N together. This Riemann surface again becomes a complex plane, a

manifold $M \infty N$ topologically $\sim S^2$. This gluing also induces conformal mappings $\hat{h}_{A_i}(w_i)$ and $\hat{h}_{B_j}(w_j)$ of the unit disks $|w_i| \leq 1$ and $|w_j| \leq 1$ of $n + m$ strings $\{A_i\}, \{B_j\}$ into the plane $M \infty N$, which are again analytic and invertible inside each disk.

Now we can state the GGRT, which was first proved by LPP²⁾ aside from the sign factor:

Theorem[LPP] *Let $\langle v(\{B_j\}, \{A_i\}) |$ be the LPP vertex defined by this set of mappings,*

$$\langle v(\{B_j\}, \{A_i\}) | \prod_{i=1}^n |A_i\rangle_{A_i} \prod_{j=1}^m |B_j\rangle_{B_j} = \left\langle \prod_{i=1}^n \hat{h}_{A_i}[\mathcal{O}_{A_i}] \prod_{j=1}^m \hat{h}_{B_j}[\mathcal{O}_{B_j}] \right\rangle_{M \infty N} . \quad (2.13)$$

Then, if the conformal anomaly is zero, the fused vertex (2.12) is just equals this LPP vertex up to a sign factor $\epsilon(A)$:

$$\langle v(D, \{B_j\}) | \langle v(C, \{A_i\}) | R(C, D) \rangle = \epsilon(A) \langle v(\{B_j\}, \{A_i\}) | . \quad (2.14)$$

The sign factor $\epsilon(A)$ is an operator

$$\epsilon(A) = (-1)^{\sum_i |A_i|} \quad (2.15)$$

which changes sign depending on the states $|A_i\rangle = \mathcal{O}_{A_i}|0\rangle$ to be contracted, where $|A_i|$ denotes statistics index defined to be 0 (1) when the operator $\mathcal{O}_{A_i}$ is bosonic (fermionic). If the conformal anomaly is present, the equality (2.14) is violated by a multiplicative c -number factor which depends non-trivially on the mappings g , h_D and h_C .

LPP analyzed the above gluing procedure in Eq. (2.12) more carefully as shown in Fig. 1. First, the complex planes M and N defining the vertices $\langle v(C, \{A_i\}) |$ and $\langle v(D, \{B_j\}) |$ are mapped by h_C^{-1} and $I \circ h_D^{-1}$ so that the exterior region of string C in M is mapped to the region outside a unit circle and the exterior region of string D in N to the inside of a unit circle, respectively. Then the region outside the unit circle in the plane $h_C^{-1}(M)$ and the region inside the unit circle in the plane $I \circ h_D^{-1}(N)$ are glued smoothly as they stand. But, unless the mappings h_C and h_D are $SL(2; C)$ transformations, neither they nor their inverses will be one-to-one mappings outside the unit circles. So the glued surface $h_C^{-1}(M) \infty I \circ h_D^{-1}(N)$ will generally possess branch-cut singularities. Since the covering surface nevertheless has a topology of S^2 , there exists a mapping g which carries the surface into the plane $M \infty N$, smoothing out the branch cuts. Therefore the conformal mappings $\hat{h}_{A_i}$ and $\hat{h}_{B_j}$ of the strings $\{A_i\}, \{B_j\}$ into the plane $M \infty N$, mentioned in the Theorem, can thus be identified with

$$\hat{h}_{A_i} = g \circ h_C^{-1} \circ h_{A_i} , \quad \hat{h}_{B_j} = g \circ I \circ h_D^{-1} \circ h_{B_j} . \quad (2.16)$$

The last step mapping g corresponds to a resmoothing procedure, explaining the name ‘Gluing and Resmoothing Theorem’.




Fig. 1. Gluing and subsequent smoothing in the contraction of two vertices.

§3. A simple proof for the tree level GGRT

We give in this section a proof for the GGRT at tree level. This proof is much simpler than the original one by LPP and, therefore, makes it easy to trace correctly the appearing sign factors.

As promised, we first discuss the parameterization forms for the conformal transformation operator U_f introduced in Eq. (2.3), and its ambiguity of multiplicative constant which exists if *the conformal anomaly (central charge c) is nonzero*. As in the usual Lie group, there are a variety of ways of representing the group elements U_f in terms of the Virasoro generators L_n . We call the parameterization form

$$U_f = \exp\left(\sum_n v_{-n} L_n\right) \quad (3.1)$$

already cited in Eq. (2.5) ‘canonical form’, which is most commonly used in the Lie group theory. Another useful parameterization form which we refer to as ‘normal ordered’ form, is

given by

$$U_f = \exp\left(\sum_{n \geq 2} v_n L_{-n}\right) \exp\left(\sum_{k=0, \pm 1} v_{-k} L_k\right) \exp\left(\sum_{m \geq 2} v_{-m} L_m\right). \quad (3.2)$$

Note that the middle factor $\exp\left(\sum_{k=0, \pm 1} v_{-k} L_k\right)$ is the element belonging to the $SL(2; C)$ subgroup. Of course we can convert various parameterization forms from one to another by using the commutation relations of L_n . But the point is that, if the conformal anomaly is nonzero, there appears a non-trivial multiplicative c-number factor in front in this rewriting; for instance, we have a relation like

$$\exp\left(\sum_{n \geq 2} v_n L_{-n}\right) \exp\left(\sum_{k=0, \pm 1} v_{-k} L_k\right) \exp\left(\sum_{m \geq 2} v_{-m} L_m\right) = e^a \exp\left(\sum_n v'_n L_n\right). \quad (3.3)$$

The front c-number factor e^a depends on the central charge c (the exponent a is linear in c), but the other group element part is uniquely determined independently of c . This means that, in the presence of nonzero central charge, the conformal transformation operator U_f has an ambiguity of overall factor depending on which parameterization form is adopted in defining U_f . This is so because unit ‘operator’ 1 is also one of the generators of the extended Virasoro algebra with central charge.

The same problem of multiplicative c-number factor arises also in the composition law of two group elements. Whatever parameterization convention is adopted for U_f and fixed, the multiplication of two elements U_f and U_g yields $U_{f \circ g}$ of composite mapping $f \circ g$ only up to a constant e^a :

$$U_f \cdot U_g = e^a U_{f \circ g}. \quad (3.4)$$

Again the constant a is linear in the central charge c , (and has complicated dependence both on the mappings f , g and the parameterization convention). So *the naive composition law is violated unless the conformal anomaly is zero*. This is the crucial property which gives the reason why the GGRT holds only in the critical dimension.

The characteristic feature of the normal ordered form (3.2) for U_f is that it manifestly satisfies

$$\langle 3 | U_f | 0 \rangle = \langle 0 | U_f | 3 \rangle = 1 \quad \left(\langle 3 | \equiv \langle 0 | c_{-1} c_0 c_1, \quad | 3 \rangle \equiv c_{-1} c_0 c_1 | 0 \rangle \right) \quad (3.5)$$

even when $c \neq 0$. (Recall that the $SL(2; C)$ invariant vacuum is normalized by the condition $\langle 0 | c_{-1} c_0 c_1 | 0 \rangle = \langle 3 | 0 \rangle = \langle 0 | 3 \rangle = 1$.) This property follows because: the $SL(2; C)$ invariant vacuum, either $|0\rangle$ or $\langle 0|$, is literally invariant under $SL(2; C)$ transformation $\exp\left(\sum_{k=0, \pm 1} v_{-k} L_k\right)$, the ghost-number 3 ket state $|3\rangle \equiv c_{-1} c_0 c_1 |0\rangle$ is invariant under $\exp\left(\sum_{m \geq 2} v_{-m} L_m\right)$ since $L_m |3\rangle = L_m c_{-1} c_0 c_1 |0\rangle = 0$ for $m \geq 1$, and similarly, $\langle 3|$ is invariant under $\exp\left(\sum_{n \geq 2} v_n L_{-n}\right)$ since $\langle 3| L_{-n} = \langle 0| c_{-1} c_0 c_1 L_{-n} = 0$ for $n \geq 1$.

Now we have the following simple proposition, which was essentially stated and used in LPP already:²⁾

Proposition 1 *Let $f(z)$ be a conformal mapping satisfying $f(0) = 0$. If $f(z)$ is analytic and invertible in a neighborhood of $z = 0$, then the corresponding operator U_f defined by Eqs. (2.3) and (2.4), leaves the $SL(2;C)$ ket vacuum inert up to a multiplicative constant:*

$$U_f |0\rangle = e^a |0\rangle. \quad (3.6)$$

If U_f is taken to be of the normal ordered form (3.2), or the conformal anomaly is absent, this constant e^a equals 1.

Proof) The $SL(2;C)$ ket vacuum $|0\rangle$ is characterized by the property that $\phi(z) |0\rangle$ remains regular as $z \rightarrow 0$ for any primary fields $\phi(z)$. This property implies for a primary field of dimension d , $\phi(z) = \sum_n \phi_n z^{-n-d}$, that

$$\lim_{z \rightarrow 0} \phi(z) |0\rangle = \text{regular} \quad \Leftrightarrow \quad \phi_n |0\rangle = 0 \quad \text{for } n \geq 1 - d. \quad (3.7)$$

So, if we can show that

$$\lim_{z \rightarrow 0} f[\phi(z)] |0\rangle = \text{regular}, \quad (3.8)$$

for any primary fields ϕ , then since $f[\phi(z)] = \sum_n f[\phi_n] z^{-n-d}$, we can deduce

$$f[\phi_n] |0\rangle = U_f \phi_n U_f^{-1} |0\rangle = 0 \quad \rightarrow \quad \phi_n U_f^{-1} |0\rangle = 0 \quad \text{for } n \geq 1 - d \quad (3.9)$$

This already implies that $U_f^{-1} |0\rangle$ is proportional to the vacuum $|0\rangle$, $U_f^{-1} |0\rangle = e^{-a} |0\rangle$, or equivalently, $U_f |0\rangle = e^a |0\rangle$ with some constant a . But, acting $\langle 3|$ to this relation from the left and using the normalization condition $\langle 3|0\rangle = 1$, we have

$$\langle 3| U_f |0\rangle = e^a. \quad (3.10)$$

If U_f is of the normal ordered form (3.2), or the conformal anomaly is absent, the left-hand side is 1 by Eq. (3.5), and $e^a = 1$ follows.

Thus we have now only to prove Eq. (3.8). The mapped field $f[\phi(z)]$ is explicitly given by Eq. (2.4) for primary fields and so is expanded as

$$f[\phi(z)] = (f'(z))^d \phi(f(z)) = \sum_n \phi_n \cdot (f(z))^{-n-d} (f'(z))^d. \quad (3.11)$$

By the assumption of analyticity of $f(z)$ around the origin and $f(0) = 0$, $f(z)$ behaves as $f(z) = f_1 z + O(z^2)$. Moreover, $f'(0) \neq 0$ by the assumption of invertibility of $f(z)$ around $z = 0$, and hence $f_1 \neq 0$. So, clearly, singular terms in the expansion (3.11) as $z \rightarrow 0$ are

only those with $-n-d \leq -1$, but they all vanish on the vacuum $|0\rangle$ by Eq. (3.7), $\phi_n |0\rangle = 0$ for $n \geq 1-d$. Thus the condition (3.8) actually holds. q.e.d.

We can rewrite U_f always into the normal ordered form up to a multiplicative constant. Then, let the general operator U_f of the normal ordered form (3.2) act on the ket vacuum:

$$\begin{aligned} U_f |0\rangle &= \exp\left(\sum_{n \geq 2} v_n L_{-n}\right) \exp\left(\sum_{k=0, \pm 1} v_{-k} L_k\right) \exp\left(\sum_{m \geq 2} v_{-m} L_m\right) |0\rangle \\ &= \exp\left(\sum_{n \geq 2} v_n L_{-n}\right) |0\rangle \end{aligned} \quad (3.12)$$

where use has been made of $L_n |0\rangle = 0$ for $n \geq -1$. So, if this equals $|0\rangle$, we must have $v_n = 0$ for $n \geq 2$. This implies that U_f has actually a simple form

$$U_f = \exp\left(\sum_{k=0, \pm 1} v_{-k} L_k\right) \exp\left(\sum_{m \geq 2} v_{-m} L_m\right) \quad (3.13)$$

for such $f(z)$ analytic and invertible around the origin. In the case of infinitesimal transformation, $f(z) = z + \delta z$, this is an expected result from the beginning, since L_{-n} is a generator of $\delta z = z^{-n+1}$ which is singular at $z = 0$ for $n \geq 2$.

In a similar manner, one can prove that the bra vacuum $\langle 0|$ remains intact under conformal transformations which have the same properties around the point at infinity.

Proposition 2 *Let $f(z)$ be a conformal mapping satisfying the same conditions as in Proposition 1. Then the mapping $g \equiv I \circ f \circ I$ satisfies, schematically writing, $g(\infty) = \infty$ and is analytic and invertible in a neighborhood of $z = \infty$. The corresponding operator $U_g = U_{I \circ f \circ I}$ leaves the $SL(2; \mathbb{C})$ bra vacuum inert up to a multiplicative constant:*

$$\langle 0| U_g = e^a \langle 0|. \quad (3.14)$$

If U_g is taken to be of the normal ordered form (3.2), or the conformal anomaly is absent, this constant e^a equals 1.

Alternatively, this could also be proved as follows if we use the form (3.13) for U_f and the transformation property of L_n under inversion, $I[L_n] = (-1)^n L_{-n}$:

$$\begin{aligned} U_g &= U_{I \circ f \circ I} = I \left[\exp\left(\sum_{k=0, \pm 1} v_{-k} L_k\right) \exp\left(\sum_{m \geq 2} v_{-m} L_m\right) \right] \\ &= \exp\left(\sum_{m \geq 2} (-1)^m v_{-m} L_{-m}\right) \exp\left(\sum_{k=0, \pm 1} (-1)^k v_{-k} L_{-k}\right). \end{aligned} \quad (3.15)$$

Then $\langle 0| U_g = \langle 0|$ would be clear since $\langle 0| L_{-n} = 0$ for $n \geq -1$.

Without loss of generality, we can assume both h_C and h_D to map the origin of the unit disk to the origins in M and N , respectively:

$$h_C(w=0) = 0, \quad h_D(w=0) = 0. \quad (3.16)$$

This is achieved, if necessary, by performing $SL(2; \mathbb{C})$ transformation on M and N , since the CFT correlation functions are $SL(2; \mathbb{C})$ invariant. Then, the mappings h_C^{-1} and h_D^{-1} , as well as h_C and h_D , satisfy the required properties of the propositions, and so we have

$$U_{h_C^{-1}} |0\rangle = |0\rangle, \quad \langle 0| U_{I \circ h_D^{-1} \circ I} = \langle 0|. \quad (3.17)$$

We shall frequently use a simple formula below which follows from Eq. (2.3) immediately:

$$\langle f[\mathcal{O}_1] f[\mathcal{O}_2] \cdots \rangle = \langle 0| U_f \mathcal{O}_1 \mathcal{O}_2 \cdots U_f^{-1} |0\rangle. \quad (3.18)$$

Now start the proof of GGRT, Eq. (2.14). Using the LPP mapping relations (2.16) and the formula (3.18), we first rewrite Eq. (2.13) as

$$\begin{aligned} & \langle v(\{B_j\}, \{A_i\}) | \prod_{i=1}^n |A_i\rangle_{A_i} \prod_{j=1}^m |B_j\rangle_{B_j} \\ &= \left\langle \prod_{i=1}^n \hat{h}_{A_i}[\mathcal{O}_{A_i}] \prod_{j=1}^m \hat{h}_{B_j}[\mathcal{O}_{B_j}] \right\rangle_{M \times N} \\ &= \left\langle \prod_{i=1}^n g \circ h_C^{-1} \circ h_{A_i}[\mathcal{O}_{A_i}] \prod_{j=1}^m g \circ I \circ h_D^{-1} \circ h_{B_j}[\mathcal{O}_{B_j}] \right\rangle_{M \times N} \\ &= \langle 0| U_g \prod_{i=1}^n h_C^{-1} \circ h_{A_i}[\mathcal{O}_{A_i}] \prod_{j=1}^m I \circ h_D^{-1} \circ h_{B_j}[\mathcal{O}_{B_j}] U_g^{-1} |0\rangle. \end{aligned} \quad (3.19)$$

Inserting the completeness relation (2.8) in the middle here, and applying the formula (3.18) for $f = h_C^{-1}$ and $I \circ h_D^{-1} \circ I$, we have

$$\begin{aligned} &= \sum_{\alpha} \langle 0| U_g \prod_{i=1}^n h_C^{-1} \circ h_{A_i}[\mathcal{O}_{A_i}] \mathcal{O}_{\alpha} |0\rangle \langle 0| I[\mathcal{O}_{\tilde{\alpha}}] \prod_{j=1}^m I \circ h_D^{-1} \circ h_{B_j}[\mathcal{O}_{B_j}] U_g^{-1} |0\rangle \\ &= \sum_{\alpha} \langle 0| U_g U_{h_C^{-1}} \prod_{i=1}^n h_{A_i}[\mathcal{O}_{A_i}] h_C[\mathcal{O}_{\alpha}] U_{h_C^{-1}}^{-1} |0\rangle \\ &\quad \times \langle 0| U_{I \circ h_D^{-1} \circ I} I \circ h_D[\mathcal{O}_{\tilde{\alpha}}] \prod_{j=1}^m I \circ h_{B_j}[\mathcal{O}_{B_j}] U_{I \circ h_D^{-1} \circ I}^{-1} U_g^{-1} |0\rangle. \end{aligned} \quad (3.20)$$

But, by Eq. (3.17), we can use $U_{h_C^{-1}} |0\rangle = |0\rangle$ and $\langle 0| U_{I \circ h_D^{-1} \circ I} = \langle 0|$ to get

$$\begin{aligned} &= \sum_{\alpha} \langle 0| U_g U_{h_C^{-1}} \prod_{i=1}^n h_{A_i}[\mathcal{O}_{A_i}] h_C[\mathcal{O}_{\alpha}] |0\rangle \\ &\quad \times \langle 0| I \circ h_D[\mathcal{O}_{\tilde{\alpha}}] \prod_{j=1}^m I \circ h_{B_j}[\mathcal{O}_{B_j}] U_{I \circ h_D^{-1} \circ I}^{-1} U_g^{-1} |0\rangle. \end{aligned} \quad (3.21)$$

Now suppose that the relations

$$\langle 0| U_g U_{h_C^{-1}} = \langle 0| , \quad U_{I \circ h_D^{-1} \circ I}^{-1} U_g^{-1} |0\rangle = |0\rangle , \quad (3.22)$$

hold, then we obtain

$$\begin{aligned} &= \sum_{\alpha} \langle 0| \prod_{i=1}^n h_{A_i}[\mathcal{O}_{A_i}] h_C[\mathcal{O}_{\alpha}] |0\rangle \cdot \langle 0| I \circ h_D[\mathcal{O}_{\tilde{\alpha}}] \prod_{j=1}^m I \circ h_{B_j}[\mathcal{O}_{B_j}] |0\rangle \\ &= \sum_{\alpha} \left\langle \prod_{i=1}^n h_{A_i}[\mathcal{O}_{A_i}] h_C[\mathcal{O}_{\alpha}] \right\rangle \cdot \left\langle I \circ h_D[\mathcal{O}_{\tilde{\alpha}}] \prod_{j=1}^m I \circ h_{B_j}[\mathcal{O}_{B_j}] \right\rangle \\ &= \sum_{\alpha} \left\langle \prod_{i=1}^n h_{A_i}[\mathcal{O}_{A_i}] h_C[\mathcal{O}_{\alpha}] \right\rangle \cdot \left\langle h_D[\mathcal{O}_{\tilde{\alpha}}] \prod_{j=1}^m h_{B_j}[\mathcal{O}_{B_j}] \right\rangle , \end{aligned} \quad (3.23)$$

where use has been made of the invariance of CFT correlation functions under inversion, i.e., $\langle I[\mathcal{O}] \rangle = \langle \mathcal{O} \rangle^*$. But the last expression is just identical with the definition of the vertices $\langle v(C, \{A_i\})|$ and $\langle v(D, \{B_j\})|$ in Eqs. (2.10) and (2.11). So, we obtain

$$\begin{aligned} &= \sum_{\alpha} \langle v(C, \{A_i\})| \prod_{i=1}^n |A_i\rangle_{A_i} |\alpha\rangle_C \cdot \langle v(D, \{B_j\})| |\tilde{\alpha}\rangle_D \prod_{j=1}^m |B_j\rangle_{B_j} \\ &= \sum_{\alpha} \langle v(D, \{B_j\})| \langle v(C, \{A_i\})| \prod_{i=1}^n |A_i\rangle_{A_i} |\alpha\rangle_C |\tilde{\alpha}\rangle_D \prod_{j=1}^m |B_j\rangle_{B_j} \\ &= \langle v(D, \{B_j\})| \langle v(C, \{A_i\})| \prod_{i=1}^n |A_i\rangle_{A_i} |R(C, D)\rangle \prod_{j=1}^m |B_j\rangle_{B_j} \\ &= \epsilon(A) \langle v(D, \{B_j\})| \langle v(C, \{A_i\})| |R(C, D)\rangle \prod_{i=1}^n |A_i\rangle_{A_i} \prod_{j=1}^m |B_j\rangle_{B_j} , \end{aligned} \quad (3.24)$$

where we have used the expression (2.9) for the reflector $|R(C, D)\rangle$, and $\epsilon(A)$ is the sign factor defined in Eq. (2.15) which has appeared since we have changed the order of $\prod_{i=1}^n |A_i\rangle_{A_i}$ and the reflector $|R(1, 2)\rangle$, the latter of which is Grassmann *odd*.**) We thus obtain the desired identity:

$$\langle v(\{B_j\}, \{A_i\})| = \epsilon(A) \langle v(D, \{B_j\})| \langle v(C, \{A_i\})| |R(C, D)\rangle . \quad (3.25)$$

Now it is, therefore, sufficient to prove the relations (3.22). Recall that g is a mapping which smoothes out the branch cuts generated in the plane $h_C^{-1}(M) \propto I \circ h_D^{-1}(N)$. So, although the mappings g as well as h_C^{-1} and h_D^{-1} are singular, the composite mapping $g \circ h_C^{-1}$

*) Incidentally, this invariance leads to an identity $\langle A|B\rangle = (-1)^{|A||B|} \langle B|A\rangle$, since $\langle A|B\rangle = \langle I[\mathcal{O}_A] \mathcal{O}_B \rangle = \langle \mathcal{O}_A I[\mathcal{O}_B] \rangle = (-1)^{|A||B|} \langle I[\mathcal{O}_B] \mathcal{O}_A \rangle = (-1)^{|A||B|} \langle B|A\rangle$. This implies the symmetry property of the reflector, $\langle R(1, 2)| = \langle R(2, 1)|$.

**) The Grassmann oddness comes from the fact $\langle 0| c_{-1} c_0 c_1 |0\rangle = 1$ after all, and hence $\mathcal{O}_{\alpha}$ and $\mathcal{O}_{\tilde{\alpha}}$ must have opposite statistics in order to have non-vanishing inner product $\langle \tilde{\alpha} | \alpha \rangle = 1$. For the case of closed string, however, the reflector $|R(1, 2)\rangle$ is Grassmann *even* since it is a product of two ‘open string’ reflectors corresponding to the holomorphic and anti-holomorphic modes.

from M to $M \infty N$, is analytic and invertible outside the image of the unit disk, $h_C(|w| < 1)$, in the M plane, and so is the mapping $g \circ I \circ h_D^{-1}$ from N to $M \infty N$, outside the image of the unit disk, $h_D(|w| < 1)$, in the N plane. Again, using the freedom of $SL(2; \mathbb{C})$ transformation in the $M \infty N$ plane, we can make the mapping g to satisfy

$$g \circ h_C^{-1}(\infty) = \infty, \quad g \circ I \circ h_D^{-1} \circ I(0) = 0. \quad (3.26)$$

Then, the mappings $g \circ h_C^{-1}$ and $g \circ I \circ h_D^{-1} \circ I$ are analytic and invertible in a neighborhood of $z = \infty$ and $z = 0$, respectively, and so we can apply the propositions 2 and 1 to obtain

$$\langle 0 | U_{g \circ h_C^{-1}} = \langle 0 |, \quad U_{g \circ I \circ h_D^{-1} \circ I} | 0 \rangle = | 0 \rangle. \quad (3.27)$$

These give just the desired relations (3.22), if

$$U_{g \circ h_C^{-1}} = U_g U_{h_C^{-1}}, \quad U_{g \circ I \circ h_D^{-1} \circ I} = U_g U_{I \circ h_D^{-1} \circ I}. \quad (3.28)$$

These actually hold with *weight one* when *the conformal anomaly is zero*. Other than in critical dimension, there appears very non-trivial multiplicative c-number factor by which Eq. (2.14) is violated. This finishes the proof of GGRT.

§4. GGRT at one loop

Next we prove here a generalized version of GGRT at one loop level; that is, the theorem for the fused vertex with double contractions by two reflectors, which read something like

$$\langle v(D, F, \{B_j\}) | \langle v(C, \{A_k\}, E) | | R(C, D) \rangle | R(E, F) \rangle = \epsilon_L \langle v_L(\{B_j\}, \{A_k\}) |. \quad (4.1)$$

Here the suffix L denotes the quantities at 1-loop level whose more precise definitions will be given in the course of the proof.

Using the GGRT at tree level, we can first rewrite the left-hand side into

$$\begin{aligned} & \langle v(D, F, \{B_j\}) | \langle v(C, \{A_k\}, E) | | R(C, D) \rangle | R(E, F) \rangle \\ &= \langle v(F, \{\Phi_i\}, E) | \epsilon(E + A) | R(E, F) \rangle \end{aligned} \quad (4.2)$$

with abbreviation $\{\Phi_i\}$ denoting the combined set of states $\{B_j\}$ and $\{A_k\}$, where $\langle v(F, \{\Phi_i\}, E) |$ is a tree level LPP vertex obtained by the fusion of the two vertices by a single contraction $| R(C, D) \rangle$, and $\epsilon(E + A)$ is the operator sign factor $(-1)^{|E| + \sum_k |A_k|}$ according to Eq. (2.15). Since this vertex $\langle v(F, \{\Phi_i\}, E) |$ is a tree level one, it corresponds to a plane, which we call

M , and there are mappings h_F , h_E and h_{Φ_i} which map the unit disks of the strings F , E and Φ_i into the plane M , analytic and invertible inside each unit disk, respectively:

$$\begin{aligned}
& \langle v(F, \{\Phi_i\}, E) | \epsilon(E + A) | R(E, F) \rangle \prod_i |\mathcal{O}_{\Phi_i}\rangle_{\Phi_i} \\
&= \sum_{\alpha} (-1)^{|\alpha| + \sum_k |A_k|} \langle v(F, \{\Phi_i\}, E) | |\alpha\rangle_E |\tilde{\alpha}\rangle_F \prod_i |\mathcal{O}_{\Phi_i}\rangle_{\Phi_i} \\
&= (-1)^{\sum_k |A_k|} \sum_{\alpha} (-1)^{|\alpha|} \left\langle h_E[\mathcal{O}_{\alpha}] h_F[\mathcal{O}_{\tilde{\alpha}}] \prod_{i=1}^n h_{\Phi_i}[\mathcal{O}_{\Phi_i}] \right\rangle_M. \quad (4.3)
\end{aligned}$$

Using the freedom of $SL(2; \mathbb{C})$, we can assume without loss of generality that

$$h_E(w=0) = 0, \quad h_F(w=0) = \infty. \quad (4.4)$$

These mappings are schematically shown in Fig. 2. The further contraction by $|R(E, F)\rangle$ in Eq. (4.2), or summation over α in Eq. (4.3), corresponds to the gluing of the two boundaries $h_E(|w|=1)$ and $h_F(|w|=1)$ in this plane M , which makes the plane a torus, which we call $M8$. The torus $M8$ can be represented by a complex plane with identification




Fig. 2. Mappings defining $\langle v(F, \{\Phi_i\}, E) |$.

$$z \sim q^2 z, \quad \exists q \equiv e^{i\pi\tau} \quad (4.5)$$

This means that there is a smooth mapping of M into the torus plane $M8$, and there are mappings $\hat{h}_{\Phi_i}$ of the unit disks to $M8$ which are analytic and invertible inside each unit circle.

In string field theory, this mapping can be decomposed into the following steps in a very similar manner to LPP at the tree level case. As shown in Fig. 3, the complex plane M is mapped in two ways, one by h_E^{-1} and the other by $I \circ h_F^{-1}$, so that the exterior region of string E in M is mapped to the region outside a unit circle and the exterior region of string F in M to the inside of a unit circle, respectively. Then the region outside the unit circle in the plane $h_E^{-1}(M)$ and the region inside the unit circle in the plane $I \circ h_F^{-1}(M)$ are glued smoothly as they stand. But, again, the glued surface generally possesses branch cuts unless the mappings h_C and h_D are $SL(2; \mathbb{C})$ transformations. Since we know the glued surface is a covering space of a torus in any case, there exists a mapping g which carries the surface into the torus plane $M8$ (the plane with identification $z \sim q^2 z$), smoothing out the branch cuts. Therefore the conformal mappings $\hat{h}_{\Phi_i}$ of the strings $\{\Phi_i\}$ into the torus plane $M8$,




Fig. 3. Gluing and resmoothing procedure to obtain the loop vertex $\langle v_L(\{\Phi_i\}) \rangle$.

mentioned above, are identified with

$$\hat{h}_{\Phi_i} = g \circ h_E^{-1} \circ h_{\Phi_i}. \quad (4.6)$$

But, in this loop case, the mappings via the other route, $g \circ I \circ h_F^{-1} \circ h_{\Phi_i}$, should equally be good mappings. Indeed, the whole region R in M outside the both images of the unit disks of strings E and F , is mapped to the two adjacent regions in $M8$ displaced with a period q^2 if we follow the two routes of mappings, $g \circ h_E^{-1}$ and $g \circ I \circ h_F^{-1}$. This can be easily seen by inspecting Fig. 3. That is, we have the following equation for $\forall z \in R$ in M

$$g \circ I \circ h_F^{-1}(z) = q^2 \times g \circ h_E^{-1}(z). \quad (4.7)$$

This is a key relation in this one-loop case. This is purely a c-number relation between the two conformal mappings. The corresponding operator relation of course reads

$$U_{g \circ I \circ h_F^{-1}} = U_{q^2 \circ g \circ h_E^{-1}}, \quad (4.8)$$

where q^2 as a mapping denotes $q^2(z) = q^2 \cdot z$. We know that the operator representation of this Weyl transformation q^2 is given by $U_{q^2} = q^{2L_0}$. Therefore, if the conformal anomaly is zero, we can use the composition law for the group elements freely to obtain

$$\begin{aligned} U_{g \circ I \circ h_F^{-1}} &= q^{2L_0} U_{g \circ h_E^{-1}} \\ \rightarrow U_{h_F \circ I}^{-1} &= U_{I \circ h_F^{-1}} = U_g^{-1} q^{2L_0} U_{g \circ h_E^{-1}}. \end{aligned} \quad (4.9)$$

If $c \neq 0$, there will appear non-trivial multiplicative c-number factors in these equations.

With these equations at hand, we can now prove the GGRT at one-loop as follows. We rewrite Eq. (4.3), aside from the fixed sign factor $(-1)^{\sum_k |A_k|}$, into

$$\begin{aligned} &\sum_{\alpha} (-1)^{|\alpha|} \left\langle h_E[\mathcal{O}_{\alpha}] h_F[\mathcal{O}_{\tilde{\alpha}}] \prod_{i=1}^n h_{\Phi_i}[\mathcal{O}_{\Phi_i}] \right\rangle_M \\ &= \sum_{\alpha} (-1)^{|\alpha|} \epsilon' \left\langle h_F[\mathcal{O}_{\tilde{\alpha}}] \prod_{i=1}^n h_{\Phi_i}[\mathcal{O}_{\Phi_i}] h_E[\mathcal{O}_{\alpha}] \right\rangle_M \\ &= \sum_{\alpha} (-1)^{|\alpha|} \epsilon' \langle 0 | U_{h_F \circ I} I[\mathcal{O}_{\tilde{\alpha}}] U_{h_F \circ I}^{-1} \prod_{i=1}^n h_{\Phi_i}[\mathcal{O}_{\Phi_i}] h_E[\mathcal{O}_{\alpha}] | 0 \rangle \end{aligned} \quad (4.10)$$

with a new sign factor $\epsilon' = (-1)^{|\alpha|(|\tilde{\alpha}| + \sum_i |\Phi_i|)}$. Fortunately, however, this sign factor is 1 since $|\alpha| + |\tilde{\alpha}| + \sum_i |\Phi_i| = 1 \pmod{2}$ in order for the correlation function in $M \sim S^2$ to be non-zero, implying $|\alpha|(|\tilde{\alpha}| + \sum_i |\Phi_i|) = |\alpha|(|\alpha| + 1) = 0 \pmod{2}$. Here, since $h_F \circ I(\infty) = \infty$ and $h_F \circ I$ is analytic and invertible in a neighborhood of $z = \infty$, we can use $\langle 0 | U_{h_F \circ I} = \langle 0 |$ by Proposition 2. Moreover we can use the above key relation (4.9) when $c = 0$. Then we further proceed as follows:

$$\begin{aligned} &= \sum_{\alpha} (-1)^{|\alpha|} \langle 0 | I[\mathcal{O}_{\tilde{\alpha}}] U_g^{-1} q^{2L_0} U_{g \circ h_E^{-1}} \prod_{i=1}^n h_{\Phi_i}[\mathcal{O}_{\Phi_i}] h_E[\mathcal{O}_{\alpha}] | 0 \rangle \\ &= \sum_{\alpha} (-1)^{|\alpha|} \langle 0 | I[\mathcal{O}_{\tilde{\alpha}}] U_g^{-1} q^{2L_0} U_{g \circ h_E^{-1}} U_{h_E} \prod_{i=1}^n h_E^{-1} \circ h_{\Phi_i}[\mathcal{O}_{\Phi_i}] \mathcal{O}_{\alpha} U_{h_E}^{-1} | 0 \rangle \\ &= \sum_{\alpha} (-1)^{|\alpha|} \langle 0 | I[\mathcal{O}_{\tilde{\alpha}}] U_g^{-1} q^{2L_0} U_g \prod_{i=1}^n h_E^{-1} \circ h_{\Phi_i}[\mathcal{O}_{\Phi_i}] \mathcal{O}_{\alpha} | 0 \rangle \\ &= \text{sTr} \left[U_g^{-1} q^{2L_0} U_g \prod_{i=1}^n h_E^{-1} \circ h_{\Phi_i}[\mathcal{O}_{\Phi_i}] \right]. \end{aligned} \quad (4.11)$$

Here, in going to the third line, we have used $U_{h_E}^{-1} | 0 \rangle = | 0 \rangle$ by Proposition 1 and $U_{g \circ h_E^{-1}} U_{h_E} = U_g$ which holds again when $c = 0$. To the last expression we have used a *definition*

$$\text{sTr}[\dots] = \sum_{\alpha} (-1)^{|\alpha|} \langle 0 | I[\mathcal{O}_{\tilde{\alpha}}] \dots \mathcal{O}_{\alpha} | 0 \rangle = \sum_{\alpha} (-1)^{|\alpha|} \langle \tilde{\alpha} | \dots | \alpha \rangle. \quad (4.12)$$

It may have sounded strange to call this a ‘definition’: of course this sTr is just the usual trace for the bosonic mode sector and the usual super trace for the ghost non-zero mode

sector. However the trace operation for the ghost zero-mode sector is not so self-evident (as will be explained later) and this gives the definition for it. The usual cyclic identity for bosonic operators also holds for this trace.

It is now immediate to rewrite the last trace expression into the final form:

$$\begin{aligned}
&= \text{sTr} \left[U_g^{-1} q^{2L_0} U_g \prod_{i=1}^n h_E^{-1} \circ h_{\Phi_i} [\mathcal{O}_{\Phi_i}] \right] = \text{sTr} \left[q^{2L_0} U_g \prod_{i=1}^n h_E^{-1} \circ h_{\Phi_i} [\mathcal{O}_{\Phi_i}] U_g^{-1} \right] \\
&= \text{sTr} \left[q^{2L_0} \prod_{i=1}^n g \circ h_E^{-1} \circ h_{\Phi_i} [\mathcal{O}_{\Phi_i}] \right] = \text{sTr} \left[q^{2L_0} \prod_{i=1}^n \hat{h}_{\Phi_i} [\mathcal{O}_{\Phi_i}] \right], \tag{4.13}
\end{aligned}$$

where the mapping relation (4.6) has been used.

Thus, if we define the CFT correlation function on the torus by

$$\langle \mathcal{O}_1 \mathcal{O}_2 \cdots \rangle_{\text{torus } \tau} = \text{sTr} [q^{2L_0} \mathcal{O}_1 \mathcal{O}_2 \cdots], \tag{4.14}$$

and the LPP vertex at one-loop level by

$$\langle v_L(\{\Phi_i\}; \tau) | \prod_i |\Phi_i\rangle_{\Phi_i} = \left\langle \prod_i \hat{h}_{\Phi_i} [\Phi_i] \right\rangle_{\text{torus } \tau}, \tag{4.15}$$

then, what we have proved is summarized in the following GGRT at one-loop, by Eqs. (4.2), (4.3), (4.10) and (4.13).

Theorem *When the conformal anomaly is zero, the fused vertex obtained by twice contractions by two reflectors, equals the LPP vertex at one-loop level up to sign:*

$$\langle v(D, F, \{B_j\}) | \langle v(C, \{A_k\}, E) | |R(C, D)\rangle |R(E, F)\rangle = \epsilon(A) \langle v_L(\{B_j\}, \{A_k\}; \tau) |, \tag{4.16}$$

where the operator sign factor $\epsilon(A)$ is given by $\epsilon(A) = (-1)^{\sum_k |A_k|}$.

A few remarks may be in order here:

If we define correlation functions on a torus for a system possessing non-zero central charge c , it is known better to replace q^{2L_0} in Eq. (4.14) with $q^{2(L_0 - c/24)}$:

$$\langle \mathcal{O}_1 \mathcal{O}_2 \cdots \rangle_{\text{torus } \tau} = \text{sTr} [q^{2(L_0 - c/24)} \mathcal{O}_1 \mathcal{O}_2 \cdots]. \tag{4.17}$$

This operator $L_0 - c/24$ can be identified with $(L_0)_{\text{cylinder}}$ on the cylinder (ρ -plane), and $-c/24$ comes from the Schwarzian derivative term in the anomalous transformation law of the energy momentum tensor under the coordinate change $\rho \rightarrow z = e^\rho$. Indeed, with this factor $(q^2)^{-c/24}$, the vacuum functional (partition function) becomes invariant under the modular transformation $\tau \rightarrow -1/\tau$ and the other correlation functions also turn to have more natural modular transformation properties. However note that this ‘improvement’ does

not cure at all the violation of the above GGRT when the net central charge is non-zero; the multiplicative c-number factor violating the theorem is a number which depends on all the details of the mappings and moduli parameters which cannot be cancelled by such a simple factor like $(q^2)^{-c/24}$. If the net central charge is zero, then one may of course calculate the correlation functions for each sectors possessing central charges separately, by using the improved definition Eq. (4.17).

It is interesting that the super trace formula Eq. (4.12) appeared automatically in our derivation. It is by no means a priori clear how the trace should be defined for the ghost zero-mode sector, since it has a off-diagonal metric structure:

$$\langle 2|1\rangle = \langle 1|2\rangle = \langle \Omega|c_0|\Omega\rangle = 1, \quad (4.18)$$

where $|1\rangle \equiv c_1|0\rangle \equiv |\Omega\rangle$ and $\langle 1| \equiv \langle 0|c_{-1} \equiv \langle \Omega|$ are ket and bra Fock vacua, and $|2\rangle \equiv c_0|\Omega\rangle$ and $\langle 2| \equiv \langle \Omega|c_0$. If we follow the definition (4.12) of supertrace $s\text{Tr}$, then noting that $|\alpha| = 1$ and 0 for $|\alpha\rangle = |1\rangle$ and $|2\rangle$, respectively, the trace in the ghost zero-mode sector, denoted by $s\text{Tr}_0$, is calculated as

$$\begin{aligned} s\text{Tr}_0[1] &= (-1)^1 \langle 2|1|1\rangle + (-1)^0 \langle 1|1|2\rangle = -1 + 1 = 0, \\ s\text{Tr}_0[c_0] &= (-1)^1 \langle 2|c_0|1\rangle + (-1)^0 \langle 1|c_0|2\rangle = 0 + 0 = 0, \\ s\text{Tr}_0[b_0] &= (-1)^1 \langle 2|b_0|1\rangle + (-1)^0 \langle 1|b_0|2\rangle = 0 + 0 = 0, \\ s\text{Tr}_0[c_0b_0] &= (-1)^1 \langle 2|c_0b_0|1\rangle + (-1)^0 \langle 1|c_0b_0|2\rangle = 0 + 1 = 1, \end{aligned} \quad (4.19)$$

and $s\text{Tr}_0[b_0c_0] = s\text{Tr}_0[1 - c_0b_0] = -1$, of course. These equations precisely show that the ghost correlation functions on the torus vanishes unless both c_0 and b_0 modes appear at least once, in conformity with the fact that there is one zero mode each for $c(z)$ and $b(z)$ in the torus case (that is, a conformal Killing vector and a holomorphic quadratic differential, respectively).

It may be noted that the supertrace can also be rewritten into a form as given by Friedan et al,⁸⁾

$$s\text{Tr}[\dots] = -\text{Tr}[(-1)^{N_{\text{FP}}} \dots] \quad (4.20)$$

in terms of the ‘usual’ trace Tr (with understanding that $\text{Tr}_0[\mathcal{O}] \equiv \langle 2|\mathcal{O}|1\rangle + \langle 1|\mathcal{O}|2\rangle$ in the zero mode sector) and the FP ghost number defined by

$$N_{\text{FP}} = c_0b_0 + \sum_{n \geq 1} (c_{-n}b_n - b_{-n}c_n) \quad (4.21)$$

which counts the ghost number from the Fock vacuum.

The authors would like to thank K. Suehiro and N. Sasakura for helpful discussions. T. K. and T. T. are supported in part by the Grant-in-Aid for Scientific Research (#10640261)

and the Grant-in-Aid (#6844), respectively, from the Ministry of Education, Science, Sports and Culture.

References

- 1) A. LeClair, M. E. Peskin and C. R. Preitschopf, Nucl. Phys. **B317** (1989), 411.
- 2) A. LeClair, M. E. Peskin and C. R. Preitschopf, Nucl. Phys. **B317** (1989), 464.
- 3) A. A. Belavin, A. M. Polyakov and A. B. Zamolodchikov, Nucl. Phys. **B241** (1984), 333.
- 4) D. Friedan, E. Martinec and S. Shenker, Nucl. Phys. **B271** (1986), 93.
- 5) T. Kugo and T. Takahashi, Prog. Theor. Phys. **99** (1998), No.4.
- 6) T. Asakawa, T. Kugo and T. Takahashi, in preparation.
- 7) H. Sonoda, Nucl. Phys. **B311** (1988/89), 401; 417.
- 8) D. Z. Freedman, S. B. Giddings, J. A. Shapiro and C. B. Thorn, Nucl. Phys. **B298** (1988), 253.
- 9) H. Kunitomo and K. Suehiro, Nucl. Phys. **B289** (1987), 157.
- 10) L. Alvarez-Gaumé, C. Gomez, G. Moore and C. Vafa, Nucl. Phys. **B303** (1988), 411.
- 11) H. Hata, K. Itoh, T. Kugo, H. Kunitomo and K. Ogawa, Phys. Rev. **D34** (1986), 2360.